

Coherence for Categorical Operadic Theories

Miles Gould
University of Glasgow

February 2, 2008

Abstract

It has long been known that every weak monoidal category A is equivalent via monoidal functors and monoidal natural transformations to a strict monoidal category $\mathbf{st}(A)$. We generalise the definition of weak monoidal category to give a definition of weak P -category for any strongly regular (operadic) theory P , and show that every weak P -category is equivalent via P -functors and P -transformations to a strict P -category. This strictification functor is then shown to have an interesting universal property.

1 Introduction

Many definitions exist of categories with some kind of “weakened” algebraic structure, in which the defining equations hold only up to coherent isomorphism. The paradigmatic example is the theory of weak monoidal categories, as presented in [12], but there are also definitions of categories with weakened versions of the structure of groups [3], Lie algebras [2], crossed monoids [1], sets acted on by a monoid [13], rigs [10], and others. A general definition of such categories-with-structure is obviously desirable, but hard in the general case. In this paper, we restrict our attention to the case of strongly regular theories (equivalently, those given by non-symmetric operads) and present possible definitions of weak P -category and weak P -functor for any non-symmetric operad P . In support of this definition, we present a generalisation of Joyal and Street’s result from [8] that every weak monoidal category is monoidally equivalent to a strict monoidal category.

The idea is to consider the strict models of our theory as algebras for an operad, then to obtain the weak models as (strict) algebras for a weakened version of that operad (which will be a **Cat**-operad). In particular, we do not make use of the pseudo-algebras of Blackwell, Kelly and Power (for which see [4]). We weaken the operad using a similar approach to that used in Penon’s definition of n -category: see [14], or [5] for a non-rigorous summary. The weak P -categories obtained are the “unbiased” ones: for instance, if P is the terminal operad (whose strict algebras are monoids), then the weak P -categories will have tensor products of all arities, not just 0 and 2.

In section 2, we present our definitions of weak P -category and weak P -functor. In section 3 we extend Joyal and Street’s proof (or rather, Leinster’s unbiased version) to the more general case of weak P -categories. In section 4, we examine the strictification functor defined in section 3, and show that it has an interesting universal property. In section 5, we explain why our approach cannot be straightforwardly extended to theories which are not given by operads, and outline some of the approaches that could be taken to deal with this.

2 Weak P -categories

By a **plain** operad, we mean what is elsewhere called a “non-symmetric” or “non- Σ ” operad, that is one with no symmetric group action defined on it. Unless stated otherwise, all our operads are plain. The category of plain operads and their morphisms is **Operad**. We are also interested in plain operads enriched in **Cat**: a **Cat-operad** is a sequence of categories $Q(0), Q(1), \dots$, a family of composition functors $\circ : Q(n) \times Q(k_1) \times \dots \times Q(k_n) \rightarrow Q(\sum k_i)$ and an identity $1_Q \in Q(1)$, satisfying the usual associativity and unit axioms (as given, for instance, in [11] section 2.2). Since operads can be thought of as one-object multicategories, we shall refer to the objects of the categories $Q(i)$ as **1-cells** and the arrows of these categories as **2-cells** of Q . **Cat-operads** and their morphisms form a category: we call this **Cat-Operad**. We could consider operads enriched in any symmetric monoidal category $\mathcal{V}$, but here we are only concerned with the cases $\mathcal{V} = \mathbf{Cat}$ or **Set**.

Let $\mathcal{V} = \mathbf{Cat}$ or **Set**, and Q be a $\mathcal{V}$ -operad. If $A \in \mathcal{V}$, we shall write $Q \circ A$ for the coproduct $\coprod_{n \in \mathbb{N}} Q(n) \times A^n$ (this notation was introduced by Kelly in [9] for clubs). An **algebra** for a $\mathcal{V}$ -operad Q is an object $A \in \mathcal{V}$ and an arrow $h : Q \circ A \rightarrow A$ which commutes with composition in Q and such that $h(1_Q, -)$ is the identity on A .

Throughout, let P be a plain (**Set**-)operad.

A **strongly regular** algebraic theory is one that can be presented using equations that use the same variables in the same order on both sides, with each variable appearing only once on each side. For instance, the theory of monoids is strongly regular, as is the theory of sets acted on by a given monoid M . The theory of commutative monoids is not strongly regular (intuitively, because of the equation $a \cdot b = b \cdot a$) and the theory of groups is not strongly regular (again intuitively, because of the equation $g \cdot g^{-1} = 1$). It can be shown, for instance as in [11] section C.1, that the strongly regular theories are exactly those given by plain operads, in the sense that the models of a strongly regular theory T are exactly the algebras for a plain operad P_T .

Plain operads are algebras for a straightforward multi-sorted algebraic theory, so (by standard arguments from universal algebra) there is an adjunction

$$\mathbf{Set}^{\mathbb{N}} \begin{array}{c} \xrightarrow{F} \\ \perp \\ \xleftarrow{U} \end{array} \mathbf{Operad}$$

The left adjoint is given by taking labelled trees, as described in [11] section 3.2. Let $D : \mathbf{Operad} \rightarrow \mathbf{Cat}\text{-}\mathbf{Operad}$ be the functor which takes discrete categories aritywise; i.e., $DP(n)$ is the discrete category on the set $P(n)$.

Definition 2.1. The **weakening of P** , $\mathbf{Wk}(P)$, is the $\mathbf{Cat}$ -operad with the same 1-cells as FUP , and the unique 2-cell structure such that the extension of the counit is a map of $\mathbf{Cat}$ -operads and is full and faithful aritywise.

More concretely, take FUP , and, for any $A, B \in FUP(n)$, place an arrow $A \rightarrow B$ iff $\varepsilon(A) = \varepsilon(B)$ (where ε is the counit of the adjunction $F \dashv U$). The composite of two arrows $A \rightarrow B \rightarrow C$ is the unique arrow $A \rightarrow C$. In particular, the arrows $A \rightarrow B$ and $B \rightarrow A$ are inverses. See Fig. 1.




Figure 1: Part of $\mathbf{Wk}(P)(3)$ with $P = 1$

This leads us immediately to the following definition:

Definition 2.2. A **weak P -category** is an algebra for $\mathbf{Wk}(P)$.

In the case $P = 1$, this reduces exactly to Leinster's definition of unbiased monoidal category in [11] section 3.1. There, two 1-cells ϕ and ψ have the same image under ε iff they have the same arity, so the categories $\mathbf{Wk}(1)(i)$ are indiscrete. We refer to the image under h of a map $q \rightarrow q'$ in $\mathbf{Wk}(P)$ as $\delta_{q,q'}$. This is clearly a natural transformation $h(q, -) \rightarrow h(q', -)$. As a special case, we write δ_q for $\delta_{q, \varepsilon(q)}$.

It is not currently clear how this definition of weak P -category relates to that of Hu and Kriz (as introduced in [7], and elucidated in Fiore's paper [6]), as I have only recently become aware of their work.

Definition 2.3. A **strict P -category** is an algebra for DP , or equivalently a weak P -category in which every component of δ is an identity arrow.

Definition 2.4. Let (A, h) and (B, h') be weak P -categories. A **weak P -functor** from (A, h) to (B, h') is a pair (G, ψ) , where $G : A \rightarrow B$ is a functor

and ψ is a sequence of natural isomorphisms $\psi_i : h'_i(1 \times G^i) \rightarrow Gh_i$, such that the following diagrams commute, for all $n, k_1, \dots, k_n \in \mathbb{N}$:

$$\begin{array}{ccc}
 & \xrightarrow{h_{k_1} \times \dots \times h_{k_n}} & \xrightarrow{h_n} \\
 1 \times 1^n \times G^{\sum k_i} \downarrow & \swarrow \psi_{k_1} \times \dots \times \psi_{k_n} & \swarrow \psi_n \downarrow G \\
 & \xrightarrow{h'_{k_1} \times \dots \times h'_{k_n}} & \xrightarrow{h'}
 \end{array}
 =
 \begin{array}{ccc}
 & \xrightarrow{h_{\sum k_i}} & \\
 1 \times 1^n \times G^{\sum k_i} \downarrow & \swarrow \psi_{\sum k_i} & \downarrow G \\
 & \xrightarrow{h'_{\sum k_i}} &
 \end{array}
 \quad (1)$$

$$\begin{array}{ccc}
 Ga & \xrightarrow{\delta'_1} & h'(1_P, Ga) \\
 \downarrow 1 & & \downarrow \psi_1 \\
 Ga & \xrightarrow{G\delta_1} & Gh(1_P, a)
 \end{array}
 \quad (2)$$

This definition is a natural generalisation of the definition of weak unbiased monoidal functor given in [11] section 3.1.

Definition 2.5. Let (F, ϕ) and (G, ψ) be weak P -functors $(A, h) \rightarrow (B, h')$. A P -transformation $\sigma : (F, \phi) \rightarrow (G, \psi)$ is a natural transformation

$$\begin{array}{ccc}
 & F & \\
 A & \xrightarrow{\quad} & B \\
 & \Downarrow \sigma & \\
 & G &
 \end{array}$$

such that

$$\begin{array}{ccc}
 Wk(P) \circ A & \xrightarrow{h} & A \\
 \downarrow 1 \circ F \quad \left(\begin{array}{c} \sigma \\ \Rightarrow \end{array} \right) \downarrow 1 \circ G & \swarrow \psi & \downarrow G \\
 Wk(P) \circ B & \xrightarrow{h'} & B
 \end{array}
 =
 \begin{array}{ccc}
 Wk(P) \circ A & \xrightarrow{h} & A \\
 \downarrow 1 \circ F & \swarrow \phi & \downarrow F \quad \left(\begin{array}{c} \sigma \\ \Rightarrow \end{array} \right) \downarrow G \\
 Wk(P) \circ B & \xrightarrow{h'} & B
 \end{array}
 \quad (3)$$

Note that there is only one possible level of strictness here. There is a 2-category, **Wk- P -Cat**, whose objects are weak P -categories, whose 1-cells are weak P -functors, and whose 2-cells are P -transformations. Similarly, there is a 2-category **Str- P -Cat** of strict P -categories, strict P -functors, and P -transformations, which can be considered a sub-2-category of **Wk- P -Cat**.

Lemma 2.6. A P -transformation $\sigma : (F, \phi) \rightarrow (G, \psi)$ is invertible as a P -transformation if and only if it is invertible as a natural transformation.

Proof. “Only if” is obvious: we concentrate on “if”. It’s enough to show that σ^{-1} is a P -transformation, which is to say that

$$\begin{array}{ccc} h(q, Ga_{\bullet}) & \xrightarrow{\psi} & Gh(q, a_{\bullet}) \\ h(q, \sigma_{a_{\bullet}}^{-1}) \downarrow & & \downarrow \sigma_{h(q, a_{\bullet})}^{-1} \\ h(q, Fa_{\bullet}) & \xrightarrow{\phi} & Fh(q, a_{\bullet}) \end{array} \quad (4)$$

commutes for all $(q, a_{\bullet}) \in Wk(P) \circ A$, and this follows from the fact that $\sigma_{h(q, a_{\bullet})} \circ \phi = \psi \circ h(q, \sigma_{a_{\bullet}})$. $\square$

3 Main Theorem

Let P be a plain operad, and $Q = Wk(P)$, with $\pi : Q \rightarrow P$ the projection map. We write composition in P as $p \circ (p_1, \dots, p_n)$, and composition in Q as $q \langle q_1, \dots, q_n \rangle$. We also adopt the $\bullet$ notation from chain complexes and write, for instance, $p_{\bullet}$ for a sequence of objects in P and $p_{\bullet}^{\bullet}$ for a double sequence. Let $Q \circ A \xrightarrow{h} A$ be a weak P -category. We construct a strict P -category $\mathbf{st}(A)$ and a weak P -functor $(F, \phi) : \mathbf{st}(A) \rightarrow A$, and show that it is an equivalence of weak P -categories.

In fact, $\mathbf{st}$ is functorial, and is left adjoint to the forgetful functor $\mathbf{Str}\text{-}P\text{-Cat} \rightarrow \mathbf{Wk}\text{-}P\text{-Cat}$ (see §4). The theorem then says that the unit of this adjunction is pseudo-invertible, and that the strict P -categories and strict P -functors form a weakly coreflective sub-2-category of $\mathbf{Wk}\text{-}P\text{-Cat}$.

Definition 3.1. Let P, Q, A be as above. The **strictification** of A , $\mathbf{st}(A)$, is defined as follows:

- An object of $\mathbf{st}(A)$ is an object of $\coprod P(i) \times A^i$.
- An arrow $(p, a_{\bullet}) \rightarrow (p', a'_{\bullet})$ in $\mathbf{st}(A)$ is an arrow $h(p, a_{\bullet}) \rightarrow h(p', a'_{\bullet})$ in A . Composition and identities are as in A .

We define an action h' of Q on $\mathbf{st}(A)$ as follows:

- On objects, h' acts by $h'(q, (p, a_{\bullet})^{\bullet}) = (\pi(q \langle p^{\bullet} \rangle), a_{\bullet}^{\bullet})$.
- Let $f_i : (p_i, a_i) \rightarrow (p'_i, a'_i)$ for $i = 0, \dots, n$. Then $h'(p, f_{\bullet})$ is the composite

$$\begin{array}{ccc} h(p \circ (p_{\bullet}), a_{\bullet}) & \xrightarrow{\delta_{p \langle p_{\bullet}^{\bullet} \rangle}^{-1}} & h(p \langle p_{\bullet}^{\bullet} \rangle, a_{\bullet}) = h(p, h(p_0, a_0), \dots, h(p_n, a_n)) \\ & \xrightarrow{h(p, f_{\bullet})} & h(p, h(p'_0, a'_0), \dots, h(p'_n, a'_n)) = h(p \langle p'_{\bullet}^{\bullet} \rangle, a'_{\bullet}) \\ & \xrightarrow{\delta_{p \langle p'_{\bullet}^{\bullet} \rangle}} & h(p \circ (p'_{\bullet}), a'_{\bullet}). \end{array}$$

Lemma 3.2. $\mathbf{st}(A)$ is a strict P -category.

Proof. The identity and strictness conditions are obvious, as is the associativity of the action on objects. We must show that the action on arrows is associative. Let $f_i^j : (p_i^j, a_{i\bullet}^j) \rightarrow (q_i^j, b_{i\bullet}^j)$, $\sigma \in Q(n)$, and $\tau_i \in Q(k_i)$ for $j = 1, \dots, k_i$ and $i = 1, \dots, n$. We wish to show that $h'(\sigma \circ (\tau_\bullet), f_\bullet^\bullet) = h'(\sigma, h'(\tau_1, f_1^\bullet), \dots, h'(\tau_n, f_n^\bullet))$.

The LHS is

$$\begin{aligned} h(\sigma \circ (\tau_\bullet) \circ (p_\bullet^\bullet), a_\bullet^\bullet) &\xrightarrow{\delta_{\sigma \circ (\tau_i) \circ (p_\bullet^\bullet)}^{-1}} h(\sigma \circ (\tau_\bullet), h(p_1^1, a_{1\bullet}^1), \dots, h(p_n^{k_n}, a_{n\bullet}^{k_n})) \\ &\xrightarrow{h(\sigma \circ (\tau_\bullet), f_\bullet^\bullet)} h(\sigma \circ (\tau_\bullet), h(q_1^1, b_{1\bullet}^1), \dots, h(q_n^{k'_n}, b_{n\bullet}^{k'_n})) \\ &\xrightarrow{\delta_{\sigma \circ (\tau_\bullet) \circ (q_\bullet^\bullet)}} h(\sigma \circ (\tau_\bullet) \circ (q_\bullet^\bullet), b_\bullet^\bullet). \end{aligned}$$

The RHS is

$$\begin{aligned} h(\sigma \circ (\tau_\bullet) \circ (p_\bullet^\bullet), a_\bullet^\bullet) &\xrightarrow{\delta_{\sigma \circ (\tau_i \circ (p_\bullet^\bullet))}^{-1}} h(\sigma, h(\tau_1 \circ (p_1^\bullet), a_{1\bullet}^\bullet), \dots, h(\tau_n \circ (p_n^\bullet), a_{n\bullet}^\bullet)) \\ &\xrightarrow{h(\sigma, h'(\tau_\bullet, f_\bullet^\bullet))} h(\sigma, h(\tau_1 \circ (q_1^\bullet), b_{1\bullet}^\bullet), \dots, h(\tau_n \circ (q_n^\bullet), b_{n\bullet}^\bullet)) \\ &\xrightarrow{\delta_{\sigma \circ (\tau_i \circ (p_\bullet^\bullet))}} h(\sigma \circ (\tau_\bullet) \circ (q_\bullet^\bullet), b_\bullet^\bullet), \end{aligned}$$

where each $h'(\tau_i, f_i^\bullet)$ is

$$\begin{aligned} h(\tau_i \circ (p_i^\bullet), a_{i\bullet}^\bullet) &\xrightarrow{\delta_{\tau_i \circ (p_i^\bullet)}^{-1}} h(\tau_i, h(p_i^1, a_{i\bullet}^1), \dots, h(p_i^{k_i}, a_{i\bullet}^{k_i})) \\ &\xrightarrow{h(\tau_i, f_i^\bullet)} h(\tau_i, h(q_i^1, b_{i\bullet}^1), \dots, h(q_i^{k_i}, b_{i\bullet}^{k_i})) \\ &\xrightarrow{\delta_{\tau_i \circ (p_i^\bullet)}} h(\tau_i \circ (q_i^\bullet), b_{i\bullet}^\bullet). \end{aligned}$$

So the equation holds if the following diagram commutes:

$$\begin{array}{ccccc} & & h(\sigma \circ (\tau_\bullet) \circ (p_\bullet^\bullet), a_\bullet^\bullet) & & \\ & \swarrow \delta_{\sigma \circ (\tau_i) \circ (p_\bullet^\bullet)}^{-1} & \downarrow \delta_{\sigma \circ (\tau_\bullet) \circ (p_\bullet^\bullet)}^{-1} & \searrow \delta_{\sigma \circ (\tau_i \circ (p_\bullet^\bullet))}^{-1} & \\ h(\sigma \circ (\tau_\bullet), h(p_\bullet^\bullet, a_\bullet^\bullet)) & \xleftarrow{\delta_{\sigma \circ (\tau_\bullet)}} & h(\sigma, h(\tau_\bullet \circ (p_\bullet^\bullet), a_\bullet^\bullet)) & \xleftarrow{h(\sigma, \delta_{\tau_\bullet \circ (p_\bullet^\bullet)}^{-1})} & h(\sigma \circ (\tau_\bullet), h(p_\bullet^\bullet, a_\bullet^\bullet)) \\ & \downarrow h(\sigma \circ (\tau_\bullet), f_\bullet^\bullet) & \downarrow h(\sigma, h'(\tau_\bullet, f_\bullet^\bullet)) & & \downarrow h(\sigma, h'(\tau_\bullet, f_\bullet^\bullet)) \\ h(\sigma \circ (\tau_\bullet), h(q_\bullet^\bullet, b_\bullet^\bullet)) & \xleftarrow{\delta_{\sigma \circ (\tau_\bullet)}} & h(\sigma, h(\tau_\bullet \circ (q_\bullet^\bullet), b_\bullet^\bullet)) & \xrightarrow{h(\sigma, \delta_{\tau_\bullet \circ (q_\bullet^\bullet)})} & h(\sigma, h(\tau_\bullet \circ (q_\bullet^\bullet), b_\bullet^\bullet)) \\ & \swarrow \delta_{\sigma \circ (\tau_\bullet) \circ (q_\bullet^\bullet)} & \downarrow \delta_{\sigma \circ (\tau_\bullet \circ (q_\bullet^\bullet))} & \swarrow \delta_{\sigma \circ (\tau_\bullet \circ (q_\bullet^\bullet))} & \\ & & h(\sigma \circ (\tau_\bullet) \circ (q_\bullet^\bullet), b_\bullet^\bullet) & & \end{array}$$

① ②

The triangles all commute because all δ s are images of arrows in Q , and there is at most one 2-cell between any two 1-cells in Q . ② commutes by the definition of $h'(\tau_i, f_i^\bullet)$, and ① commutes by naturality of δ . $\square$

Lemma 3.3. Let $Q \circ A \xrightarrow{h} A$ and $Q \circ B \xrightarrow{h'} B$ be weak P -categories, $(F, \pi) : A \rightarrow B$ be a weak P -functor, and $(F, G, \eta, \varepsilon)$ be an adjoint equivalence. Then G naturally carries the structure of a weak P -functor, and $(F, G, \eta, \varepsilon)$ is an adjoint equivalence in **Wk-P-Cat**.

Proof. We want a sequence $(\psi_\bullet)$ of natural transformations:

$$\begin{array}{ccc} Q(i) \times B^i & \xrightarrow{h'_i} & B \\ 1 \times G^i \downarrow & \nearrow \psi_i & \downarrow G \\ Q(i) \times A^i & \xrightarrow{h_i} & A \end{array}$$

Let ψ_i be given by $Gh'(1 \times \varepsilon^i) \circ G\pi_i^{-1} \circ \eta_{h'}$, i.e.


$$\begin{array}{ccc} Q(i) \times B^i & \xrightarrow{h'_i} & B \\ 1 \times G^i \downarrow & \nearrow \psi_i & \downarrow G \\ Q(i) \times A^i & \xrightarrow{h_i} & A \end{array} = \begin{array}{ccccc} Q(i) \times B^i & \xrightarrow{1} & Q(i) \times B^i & \xrightarrow{h'_i} & B \\ 1 \times G^i \downarrow & \nearrow 1 \times \varepsilon^i & \nearrow 1 \times F^i & \nearrow \pi_i^{-1} & \nearrow F \\ Q(i) \times A^i & \xrightarrow{h_i} & A & \xrightarrow{1} & A \end{array}$$

We must check that ψ satisfies (1) and (2). For (1):

$$\begin{aligned} \text{LHS} &= \begin{array}{ccc} h'_{k_1} \times \dots \times h'_{k_n} & \xrightarrow{\quad} & h'_n \\ \downarrow 1 \times G^{\Sigma k_i} & \nearrow \psi_{k_1} \times \dots \times \psi_{k_n} & \downarrow 1 \times G^n \\ h_{k_1} \times \dots \times h_{k_n} & \xrightarrow{\quad} & h \end{array} \\ &= \begin{array}{ccccccc} & 1 & & h'_{k_1} \times \dots \times h'_{k_n} & & 1 & & h'_n \\ \downarrow 1 \times G^{\Sigma k_i} & \nearrow 1 \times \varepsilon^{\Sigma k_i} & \nearrow \pi_{k_1}^{-1} \times \dots \times \pi_{k_n}^{-1} & \nearrow 1 \times F^n & \nearrow 1 \times G^n & \nearrow 1 \times F^n & \nearrow \pi_n^{-1} & \nearrow F \\ h_{k_1} \times \dots \times h_{k_n} & \xrightarrow{\quad} & 1 & \xrightarrow{\quad} & h & \xrightarrow{\quad} & 1 & \xrightarrow{\quad} \end{array} \end{aligned}$$

$$\begin{aligned}
&= \begin{array}{ccccc}
& & 1 & \xrightarrow{\quad} & h'_{k_1} \times \cdots \times h'_{k_n} & \xrightarrow{\quad} & 1 & \xrightarrow{\quad} & h'_n & \xrightarrow{\quad} & \\
1 \times G^{\sum k_i} \downarrow & \nearrow 1 \times \varepsilon^{\sum k_i} & & \nearrow \pi_{k_1}^{-1} \times \cdots \times \pi_{k_n}^{-1} & & \nearrow 1 \times F^n & & \nearrow \pi_n^{-1} & & \nearrow F & & \downarrow G \\
& 1 \times F^{\sum k_i} & \uparrow & & 1 \times F^n & = & 1 \times F^n & & & & & \\
& h_{k_1} \times \cdots \times h_{k_n} & \xrightarrow{\quad} & 1 & \xrightarrow{\quad} & h & \xrightarrow{\quad} & 1 & \xrightarrow{\quad} & & &
\end{array} \\
&= \begin{array}{ccccc}
& & 1 & \xrightarrow{\quad} & h'_{k_1} \times \cdots \times h'_{k_n} & \xrightarrow{\quad} & h'_n & \xrightarrow{\quad} & \\
1 \times G^{\sum k_i} \downarrow & \nearrow 1 \times \varepsilon^{\sum k_i} & & \nearrow \pi_{k_1}^{-1} \times \cdots \times \pi_{k_n}^{-1} & & \nearrow 1 \times F^n & & \nearrow \pi_n^{-1} & & \nearrow F & & \downarrow G \\
& 1 \times F^{\sum k_i} & \uparrow & & 1 \times F^n & & & & & & & \\
& h_{k_1} \times \cdots \times h_{k_n} & \xrightarrow{\quad} & h & \xrightarrow{\quad} & 1 & \xrightarrow{\quad} & & & & &
\end{array} \\
&= \begin{array}{ccccc}
& & 1 & \xrightarrow{\quad} & h'_{\sum k_i} & \xrightarrow{\quad} & \\
1 \times G^{\sum k_i} \downarrow & \nearrow 1 \times \varepsilon^{\sum k_i} & & \nearrow \pi_{\sum k_i}^{-1} & & \nearrow F & & \downarrow G \\
& 1 \times F^{\sum k_i} & \uparrow & & & & & \\
& h_{\sum k_i} & \xrightarrow{\quad} & 1 & \xrightarrow{\quad} & & &
\end{array} \\
&= \begin{array}{ccc}
& h'_{\sum k_i} & \\
1 \times G^{\sum k_i} \downarrow & \nearrow \psi_{\sum k_i} & \downarrow G \\
& h_{\sum k_i} &
\end{array} \\
&= \text{RHS.}
\end{aligned}$$

For (2), consider the following diagram:



(2) is the outside of the diagram. ① commutes by the triangle identities. ② commutes by naturality of η . ③ commutes since (F, π) is a P -functor. ④ commutes by naturality of δ . ⑤ is the definition of ψ . Hence the whole diagram commutes, and (G, ψ) is a P -functor.

To see that $(F, G, \eta, \varepsilon)$ is a P -equivalence, it is now enough to show that η and ε are P -transformations, since they satisfy the triangle identities by hypothesis.


Write $(GF, \chi) = (G, \psi) \circ (F, \pi)$. We wish to show that η is a P -transformation $(1, 1) \rightarrow (GF, \chi)$. Each $\chi_{q, a_\bullet}$ is the composite

$$h(q, GFa_\bullet) \xrightarrow{\psi} Gh(q, Fa_\bullet) \xrightarrow{G\pi} GFh(q, a_\bullet)$$

Applying the definition of ψ , this is

$$h(q, GFa_\bullet) \xrightarrow{\eta} GFh(q, GFa_\bullet) \xrightarrow{G\pi^{-1}} Gh(q, FGFa_\bullet) \xrightarrow{Gh_q \varepsilon F} Gh(q, Fa_\bullet) \xrightarrow{G\pi} GFh(q, a_\bullet)$$

The axiom on η is the outside of the diagram



① commutes by naturality of η , ② commutes by naturality of π^{-1} , and ③ commutes since $G\pi \circ G\pi^{-1} = G(\pi \circ \pi^{-1}) = G1 = 1G$. The triangle commutes by the triangle identities. So the whole diagram commutes, and η is a P -transformation. By Lemma 2.6, η^{-1} is also a P -transformation. Similarly, ε and ε^{-1} are P -transformations.

□

Theorem 3.4. *Let $Q \circ A \xrightarrow{h} A$ be a weak P -category. Then A is equivalent to $\mathbf{st}(A)$ via weak P -functors and P -transformations.*

Proof. Let $F : \mathbf{st}(A) \rightarrow A$ be given by $F(p, a_\bullet) = h(p, a_\bullet)$ and identification of maps. This is certainly full and faithful, and it's essentially surjective on objects because $\delta_{1_Q}^{-1} : h(1_P, a) \rightarrow a$ is an isomorphism. It remains to show that F is a weak P -functor.

We must find a sequence $(\phi_i : h_i(1 \times F^i) \rightarrow Fh')$ of natural transformations satisfying equations (1) and (2). We can take $(\phi_i)_{q, (p_\bullet, a_\bullet)} = (\delta_{q \langle p_\bullet \rangle})_{a_\bullet}$. For (1), we must show that

$$\begin{array}{ccc}
 & \xrightarrow{h'_{k_1} \times \dots \times h'_{k_n}} & \xrightarrow{h'_n} \\
 1 \times F^{\Sigma k_i} \downarrow & \begin{array}{c} \phi_{k_1} \times \dots \times \phi_{k_n} \\ \nearrow \\ 1 \times F^n \end{array} & \downarrow \phi_n \\
 & \xrightarrow{h_{k_1} \times \dots \times h_{k_n}} & \xrightarrow{h} \\
 & & \downarrow F
 \end{array} = \begin{array}{ccc}
 & \xrightarrow{h'_{\Sigma k_i}} & \\
 1 \times F^{\Sigma k_i} \downarrow & \begin{array}{c} \phi_{\Sigma k_i} \\ \nearrow \end{array} & \downarrow F \\
 & \xrightarrow{h_{\Sigma k_i}} &
 \end{array}$$

All 2-cells in this equation are instances of δ . Since there is at most one 2-cell between two 1-cells in Q , the equation holds.

For (2) to hold, we must have

$$\begin{array}{ccc}
 F(p, a_\bullet) & \xrightarrow{\delta_{1_Q}} & h(1_P, F(p, a_\bullet)) \\
 \downarrow 1 & & \downarrow \phi_{1_P} \\
 F(p, a_\bullet) & \xrightarrow{F\delta'_{1_Q}} & Fh'(1_P, (p, a_\bullet))
 \end{array} \tag{5}$$

Since $\mathbf{st}(A)$ is a strict monoidal category, $\delta' = 1$. Apply this observation, and the definitions of F , ϕ and h' ; then (5) becomes

$$\begin{array}{ccc}
 h(p, a_\bullet) & \xrightarrow{\delta_{1_Q}} & h(1_P, h(p, a_\bullet)) \\
 \downarrow 1 & & \downarrow \delta_{1_P \langle p \rangle} \\
 h(p, a_\bullet) & \xrightarrow{1} & h(p, a_\bullet)
 \end{array}$$

Since there is at most one arrow between two 1-cells in Q , this diagram commutes. So (F, ϕ) is a weak P -functor. Hence, by Lemma 3.3, A is equivalent to $\mathbf{st}(A)$ via weak P -functors and P -transformations. $\square$

4 Significance of $\mathbf{st}$

Theorem 4.1. *Let U' be the forgetful functor $\mathbf{Str}\text{-}P\text{-Cat} \rightarrow \mathbf{Wk}\text{-}P\text{-Cat}$ (considering both of these as 1-categories). Then $\mathbf{st}$ is left adjoint to U' .*

Proof. For each $A \in \mathbf{Wk}\text{-}P\text{-}\mathbf{Cat}$, we construct an initial object $A \xrightarrow{(F', \psi)} \mathbf{st}(A)$ of the comma category $(A \downarrow U')$, thus showing that $\mathbf{st}$ is functorial and that $\mathbf{st} \dashv U'$ (and that (F', ψ) is the component of the unit at A). Let (B, h'') be a strict P -category, and $(G, \gamma) : A \rightarrow U'B$ be a weak P -functor. We must show that there is a unique strict P -functor H making the following diagram commute:

$$\begin{array}{ccc} & A & \\ (F', \psi) \swarrow & & \searrow (G, \gamma) \\ U' \mathbf{st}(A) & \xrightarrow{(H, \text{id})} & U'B \end{array} \quad (6)$$

(F', ψ) is given as follows:

- If $a \in A$, then $F'(a) = (1, a)$.
- If $f : a \rightarrow a'$ in A then $F'f$ is the lifting of $h(1, f)$ with source $(1, a)$ and target $(1, a')$.
- $\psi_{(p, a_\bullet)}$ is the lifting of $(\delta_{1_Q})_{h(p, a_\bullet)} : h(p, a_\bullet) \rightarrow h(1, h(p, a_\bullet))$ to a morphism $h'(p, F'(a)_\bullet) = (p, a_\bullet) \rightarrow (1, h(p, a_\bullet)) = F'(h(p, a_\bullet))$.

For commutativity of (6), we must have $H(1, a) = G(a)$, and for strictness of H , we must have $H(p, a_\bullet) = h''(p, H(1, a)_\bullet)$. These two conditions completely define H on objects.

Now, take a morphism $f : (p, a_\bullet) \rightarrow (p', a'_\bullet)$, which is a lifting of a morphism $g : h(p, a_\bullet) \rightarrow h(p', a'_\bullet)$ in A . Hf is a morphism $h''(p, Ga_\bullet) \rightarrow h''(p', Ga'_\bullet)$: the obvious thing for it to be is the composite

$$h''(p, Ga_\bullet) \xrightarrow{\gamma} Gh''(p, a_\bullet) \xrightarrow{Gg} Gh''(p', a'_\bullet) \xrightarrow{\gamma^{-1}} h''(p', Ga'_\bullet)$$

and we shall show that this is in fact the only possibility. Consider the composite

$$(1, h(p, a_\bullet)) \xrightarrow{\psi^{-1}} (p, a_\bullet) \xrightarrow{f} (p', a'_\bullet) \xrightarrow{\psi} (1, h(p', a'_\bullet))$$

in $\mathbf{st}(A)$. Composition in $\mathbf{st}(A)$ is given by composition in A , so this is equal to the lifting of $\delta_{1_Q} \circ g \circ \delta_{1_Q}^{-1} = h(1, g)$ to a morphism $(1, h(p, a_\bullet)) \rightarrow (1, h(p', a'_\bullet))$, namely $F'g$. So $f = \psi^{-1} \circ F'g \circ \psi$, and $Hf = H\psi^{-1} \circ HF'g \circ H\psi$. By commutativity of (6), $HF' = G$ and $H\psi = \gamma$, so $Hf = \gamma^{-1} \circ Gg \circ \gamma$ as required.

This completely defines H . So we have constructed a unique H which makes (6) commute and which is strict. Hence $(F', \psi) : A \rightarrow U' \mathbf{st}(A)$ is initial in $(A \downarrow U')$, and so $\mathbf{st} \dashv U'$. $\square$

The P -functor $(F, \phi) : \mathbf{st}(A) \rightarrow A$ constructed in Theorem 3.4 is pseudo-inverse to (F', ψ) , which we have just shown to be the A -component of the unit of the adjunction $\mathbf{st} \dashv U'$. We can therefore say that $\mathbf{Str}\text{-}P\text{-}\mathbf{Cat}$ is a weakly coreflective sub-2-category of $\mathbf{Wk}\text{-}P\text{-}\mathbf{Cat}$. Note that the counit is *not* pseudo-invertible, so this is not a 2-equivalence.

5 Further Work

Very few interesting theories are strongly regular, so this definition is unsatisfactory as it stands. It can be straightforwardly extended to theories given by symmetric operads, but to deal with the interesting cases of groups, rings, Lie algebras, etc, we must either abandon operads and move to a more expressive formalism (for instance that of Lawvere theories), or extend the notion of an operad until it is sufficiently expressive. I have taken the latter approach: by allowing any function of finite sets, and not just permutations, to act on the sets $P(i)$, we obtain a notion of operad that is equivalent in power to clones or Lawvere theories (as was proved by Tronin in [15]).

However, naïvely extending definition 2.1 to these more general operads doesn't work, as weakening the theory of commutative monoids gives the theory of strictly symmetric weak monoidal categories, rather than that of symmetric weak monoidal categories as desired. I have been working off and on on various other approaches, mainly concerned with constructing $\mathbf{Wk}(P)$ using some universal property, and have obtained some interesting early results.

References

- [1] Pierre Ageron. Les catégories monoïdales croisées. 78th PSSL, Strasbourg, Feb 2002. Available at <http://www.math.unicaen.fr/~ageron/Strasbourg.dvi>.
- [2] John C. Baez and Alissa S. Crans. Higher-dimensional algebra VI: Lie 2-algebras. *Theory and Applications of Categories*, 12:492–538, 2004, math.QA/0307263.
- [3] John C. Baez and Aaron D. Lauda. Higher-dimensional algebra V: 2-groups. *Theory and Applications of Categories*, 12:423–491, 2004, math.QA/0307200.
- [4] R. Blackwell, G.M. Kelly, and A.J. Power. Two-dimensional monad theory. *Journal of Pure and Applied Algebra*, 59:1–41, 1989.
- [5] Eugenia Cheng and Aaron Lauda. Higher-dimensional categories: an illustrated guide book, Jun 2004.
- [6] Thomas M. Fiore. Pseudo limits, biadjoints, and pseudo algebras: Categorical foundations of conformal field theory. *Memoirs of the AMS*, to appear, math.CT/04028298.
- [7] P. Hu and I. Kriz. Conformal field theory and elliptic cohomology. *Advances in Mathematics*, 189(2):325–412, 2004.
- [8] André Joyal and Ross Street. Braided tensor categories. *Advances in Mathematics*, 102:20–78, 1993.

- [9] G. M. Kelly. *An Abstract Approach To Coherence*, pages 106–174. Number 281 in Lecture Notes in Mathematics. Springer-Verlag, May 1972.
- [10] Miguel L. Laplaza. *Coherence for distributivity*, pages 29–65. Number 281 in Lecture Notes in Mathematics. Springer-Verlag, May 1972.
- [11] Tom Leinster. *Higher Operads, Higher Categories*. Cambridge University Press, 2003, math.CT/0305049.
- [12] Saunders Mac Lane. *Categories for the Working Mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, second edition, 1998.
- [13] Viktor Ostrik. Module categories, weak Hopf algebras and modular invariants. *Transform. Groups*, 8(2):159–176, 2003, math.QA/0111139.
- [14] J. Penon. Approche polygraphique des ∞ -catégories non strictes. *Cahiers de Topologie et Géométrie Différentielle Catégoriques*, XL(1):31–80, 1999.
- [15] S.N. Tronin. Abstract clones and operads. *Siberian Mathematical Journal*, 43(4):746–755, Jul 2002.